

Anti-GAK antibody (70-150 N-Term) (STJ93199)

STJ93199

GENERAL INFORMATION


Product Type	Primary antibodies
Short Description	Rabbit polyclonal antibody anti-Cyclin-G-Associated Kinase (70-150 N-Term) is suitable for use in Western Blot, Immunohistochemistry, Immunofluorescence, Immunocytochemistry and ELISA research applications.
Applications	WB, IHC-P, IF, ICC, ELISA
Host/Source	Rabbit
Reactivity	Human, Mouse

PRODUCT PROPERTIES


Clonality	Polyclonal
Clone ID	
Concentration	1 mg/mL
Conjugation	Unconjugated
Purification	The antibody was affinity-purified from rabbit anti-serum by affinity-chromatography.
Dilution Range	WB 1:500-1:2000 IHC 1:100-1:300 IF 1:200-1:1000 ELISA 1:5000
Formulation	PBS, 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.
Isotype	IgG
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	2580
Gene Symbol	GAK
Uniprot ID	GAK_HUMAN
Immunogen	The antiserum was produced against synthesized peptide derived from human GAK at amino acid range 101-150
Immunogen Region	70-150 N-Term
Specificity	GAK polyclonal antibody (Cyclin-G-Associated Kinase) binds to endogenous Cyclin-G-Associated Kinase at the amino acid region 70-150 N-Term.
Immunogen Sequence	



Western blot analysis of lysates from 293 cells, using GAK Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HepG2 cells using GAK antibody.



Western blot analysis of various cells using GAK Polyclonal Antibody

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081