

Anti-MED21 antibody (40-120 Internal) (STJ95775)

STJ95775

GENERAL INFORMATION


| | |
|--------------------------|---|
| Product Type | Primary antibodies |
| Short Description | Rabbit polyclonal antibody anti-Mediator Of Rna Polymerase Ii Transcription Subunit 21 (40-120 Internal) is suitable for use in Immunohistochemistry, Immunofluorescence and ELISA research applications. |
| Applications | IHC-P, IF-P, ELISA |
| Host/Source | Rabbit |
| Reactivity | Human, Mouse |

PRODUCT PROPERTIES

| | |
|----------------------|--|
| Clonality | Polyclonal |
| Clone ID | |
| Concentration | 1 mg/mL |
| Conjugation | Unconjugated |
| Purification | The antibody was affinity-purified from rabbit anti-serum by affinity-chromatography. |
| Dilution | IHC 1:100-1:300 |
| Range | ELISA 1:20000 |
| Formulation | PBS, 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide. |
| Isotype | IgG |
| Storage | Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles. |
| Instruction | |

TARGET INFORMATION

| | |
|---------------------------|---|
| Gene ID | 9412 |
| Gene Symbol | MED21 |
| Uniprot ID | MED21_HUMAN |
| Immunogen | The antiserum was produced against synthesized peptide derived from human MED21 at amino acid range 71-120 |
| Immunogen Region | 40-120 Internal |
| Specificity | MED21 polyclonal antibody (Mediator Of Rna Polymerase Ii Transcription Subunit 21) binds to endogenous Mediator Of Rna Polymerase Ii Transcription Subunit 21 at the amino acid region 40-120 Internal. |
| Immunogen Sequence | |



Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using MED21 Antibody. The picture on the right is blocked with the synthesized peptide.