

## Anti-TNFAIP8 antibody (1-80 Internal) (STJ96050)

STJ96050

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	Rabbit polyclonal antibody anti-Tumor Necrosis Factor Alpha-Induced Protein 8 (1-80 Internal) is suitable for use in Western Blot, Immunohistochemistry, Immunofluorescence, Immunocytochemistry and ELISA research applications.
<b>Applications</b>	WB, IHC-P, IF, ICC, ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human, Mouse

### PRODUCT PROPERTIES


<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	1 mg/mL
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	The antibody was affinity-purified from rabbit anti-serum by affinity-chromatography.
<b>Dilution</b>	WB 1:500-1:2000
<b>Range</b>	IHC 1:100-1:300 IF 1:200-1:1000 ELISA 1:10000
<b>Formulation</b>	PBS, 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.
<b>Isotype</b>	IgG
<b>Storage</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
<b>Instruction</b>	

### TARGET INFORMATION


<b>Gene ID</b>	25816
<b>Gene Symbol</b>	TNFAIP8
<b>Uniprot ID</b>	TFIP8_HUMAN
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from human TFIP8 at amino acid range 31-80
<b>Immunogen Region</b>	1-80 Internal
<b>Specificity</b>	TNFAIP8 polyclonal antibody (Tumor Necrosis Factor Alpha-Induced Protein 8) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 8 at the amino acid region 1-80 Internal.
<b>Immunogen Sequence</b>	



Western blot analysis of lysates from HUVEC cells, using TFIP8 Antibody. The lane on the right is blocked with the synthesized peptide.



Immunofluorescence analysis of A549 cells, using TFIP8 Antibody. The picture on the right is blocked with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using TFIP8 Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of various cells using TNF-IP 8 Polyclonal Antibody. Secondary antibody was diluted at 1:20000

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081