

Anti-TNFAIP2 antibody (100-180 Internal) (STJ96055)

GENERAL INFORMATION

Product Type Primary antibodies

Short Rabbit polyclonal antibody anti-Tumor Necrosis Factor Alpha-Induced Protein 2 (100-180 Internal) is suitable for use in

Description Immunohistochemistry, Immunofluorescence, Immunocytochemistry and ELISA research applications.

Applications IHC-P, IF, ICC, ELISA

Host/Source Rabbit Reactivity Human, Mouse

PRODUCT PROPERTIES

Clonality Polyclonal

Clone ID

Concentration 1 mg/mL

Conjugation Unconjugated

Purification The antibody was affinity-purified from rabbit anti-serum by affinity-chromatography.

Dilution IHC 1:100-1:300 Range IF 1:200-1:1000 ELISA 1:20000

Formulation PBS, 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.

Isotype IgG

Storage Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

Instruction

TARGET INFORMATION

Gene ID 7127

Gene Symbol TNFAIP2

Uniprot ID TNAP2_HUMAN

Immunogen The antiserum was produced against synthesized peptide derived from human TNAP2 at amino acid range 131-180


Immunogen 100-180 Internal


Region

Specificity TNFAIP2 polyclonal antibody (Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endogenous Tumor Necrosis Factor Alpha-Induced Protein 2) binds to endoge

Induced Protein 2 at the amino acid region 100-180 Internal

Immunogen Sequence





Immunofluorescence analysis of HUVEC cells, using TNAP2 Antibody. The picture on the right is blocked with the synthesized peptide.