

## Anti-HGFAC antibody (C-Term) (STJ96435)

STJ96435

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	Rabbit polyclonal antibody anti-Hepatocyte Growth Factor Activator (C-Term) is suitable for use in Western Blot and ELISA research applications.
<b>Applications</b>	WB, ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human, Mouse, Rat

### PRODUCT PROPERTIES

<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	1 mg/mL
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	The antibody was affinity-purified from rabbit anti-serum by affinity-chromatography.
<b>Dilution Range</b>	WB 1:500-1:2000 ELISA 1:40000
<b>Formulation</b>	PBS, 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.
<b>Isotype</b>	IgG
<b>Storage</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
<b>Instruction</b>	

### TARGET INFORMATION

<b>Gene ID</b>	3083
<b>Gene Symbol</b>	HGFAC
<b>Uniprot ID</b>	HGFA_HUMAN
<b>Immunogen</b>	Synthesized peptide derived from the C-terminal region of human HGFA.
<b>Immunogen Region</b>	C-Term
<b>Specificity</b>	HGFAC polyclonal antibody (Hepatocyte Growth Factor Activator) binds to endogenous Hepatocyte Growth Factor Activator at the amino acid region C-Term.
<b>Immunogen Sequence</b>	



Western blot analysis of extracts from K562, Jurkat cells, using HGFA Polyclonal Antibody. Secondary antibody was diluted at 1:20000