

## Anti-FCGR2A antibody (31-80 Internal) (STJ96610)

STJ96610

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	Rabbit polyclonal antibody anti-Low Affinity Immunoglobulin Gamma Fc Region Receptor II-A (31-80 Internal) is suitable for use in Western Blot and ELISA research applications.
<b>Applications</b>	WB, ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human, Rat, Mouse

### PRODUCT PROPERTIES


<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	1 mg/mL
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	The antibody was affinity-purified from rabbit anti-serum by affinity-chromatography.
<b>Dilution</b>	WB 1:500-1:2000
<b>Range</b>	ELISA 1:20000
<b>Formulation</b>	PBS, 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.
<b>Isotype</b>	IgG
<b>Storage</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
<b>Instruction</b>	

### TARGET INFORMATION

<b>Gene ID</b>	2212
<b>Gene Symbol</b>	FCGR2A
<b>Uniprot ID</b>	FCG2A_HUMAN
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from the Internal region of human FCGR2A at amino acid range 41-90
<b>Immunogen Region</b>	31-80 Internal
<b>Specificity</b>	FCGR2A polyclonal antibody (Low Affinity Immunoglobulin Gamma Fc Region Receptor II-A) binds to endogenous Low Affinity Immunoglobulin Gamma Fc Region Receptor II-A at the amino acid region 31-80 Internal.
<b>Immunogen Sequence</b>	



Western blot analysis of lysate from K562 cells, using FCGR2A Antibody.



Western blot analysis of K562 cells using CD32-A Polyclonal Antibody. Secondary antibody was diluted at 1:20000

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
 St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081