

## Anti-AICDA antibody (81-130 Internal) (STJ97261)

STJ97261

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	Rabbit polyclonal antibody anti-Single-Stranded Dna Cytosine Deaminase (81-130 Internal) is suitable for use in Western Blot, Immunohistochemistry, Immunofluorescence and ELISA research applications.
<b>Applications</b>	WB, IHC-P, IF-P, ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human, Mouse, Rat

### PRODUCT PROPERTIES


<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	1 mg/mL
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	The antibody was affinity-purified from rabbit anti-serum by affinity-chromatography.
<b>Dilution Range</b>	WB 1:500-1:2000 IHC 1:100-1:300 ELISA 1:10000
<b>Formulation</b>	PBS, 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.
<b>Isotype</b>	IgG
<b>Storage Instruction</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

### TARGET INFORMATION

<b>Gene ID</b>	57379
<b>Gene Symbol</b>	AICDA
<b>Uniprot ID</b>	AICDA_HUMAN
<b>Immunogen</b>	The antiserum was produced against synthesized peptide derived from the Internal region of human AICDA at amino acid range 81-130
<b>Immunogen Region</b>	81-130 Internal
<b>Specificity</b>	AICDA polyclonal antibody (Single-Stranded Dna Cytosine Deaminase) binds to endogenous Single-Stranded Dna Cytosine Deaminase at the amino acid region 81-130 Internal.
<b>Immunogen Sequence</b>	



Immunohistochemical analysis of paraffin-embedded human colon cancer, antibody was diluted at 1:100



Western blot analysis of HeLa, HepG2 cells using AID Polyclonal Antibody. Secondary antibody was diluted at 1:20000

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081