

## Anti-BPTF antibody [2F10] (STJ97882)

STJ97882

### GENERAL INFORMATION



<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	Mouse monoclonal antibody anti-Nucleosome-Remodeling Factor Subunit Bptf is suitable for use in Western Blot and ELISA research applications.
<b>Applications</b>	WB, ELISA
<b>Host/Source</b>	Mouse
<b>Reactivity</b>	Human

### PRODUCT PROPERTIES

<b>Clonality</b>	Monoclonal
<b>Clone ID</b>	2F10
<b>Concentration</b>	
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	The antibody was isolated from ascitic fluid by immunoaffinity chromatography using antigens coupled to agarose beads.
<b>Dilution Range</b>	WB 1:500-1:2000 ELISA 1:10000
<b>Formulation</b>	Ascitic fluid, 0.03% Sodium Azide, 0.5% BSA, 50% Glycerol.
<b>Isotype</b>	IgG2b
<b>Storage</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
<b>Instruction</b>	

### TARGET INFORMATION

<b>Gene ID</b>	2186
<b>Gene Symbol</b>	BPTF
<b>Uniprot ID</b>	BPTF_HUMAN
<b>Immunogen</b>	Purified recombinant fragment of human BPTF expressed in E.coli.
<b>Immunogen Region</b>	
<b>Specificity</b>	BPTF monoclonal antibody (Nucleosome-Remodeling Factor Subunit Bptf) binds to endogenous Nucleosome-Remodeling Factor Subunit Bptf.
<b>Immunogen Sequence</b>	



Western blot analysis using BPTF monoclonal antibody against HEK293 (1) and BPTF (AA: 503-670) -hlgGfC transfected HEK293 (2) cell lysate.