

## Anti-FGR antibody [6G2] (STJ97953)

STJ97953

### GENERAL INFORMATION


|                          |   |
|--------------------------|---|
| <b>Product Type</b>      | Primary antibodies  |
| <b>Short Description</b> | Mouse monoclonal antibody anti-Tyrosine-Protein Kinase Fgr is suitable for use in Western Blot, Immunohistochemistry, Immunofluorescence and ELISA research applications. |
| <b>Applications</b>      | WB, IHC-P, IF-P, ELISA  |
| <b>Host/Source</b>       | Mouse   |
| <b>Reactivity</b>        | Human, Mouse  |

### PRODUCT PROPERTIES


|                            |  |
|----------------------------|--|
| <b>Clonality</b>           | Monoclonal   |
| <b>Clone ID</b>            | 6G2  |
| <b>Concentration</b>       |  |
| <b>Conjugation</b>         | Unconjugated   |
| <b>Purification</b>        | The antibody was isolated from ascitic fluid by immunoaffinity chromatography using antigens coupled to agarose beads. |
| <b>Dilution Range</b>      | WB 1:500-1:2000<br>IHC 1:200-1:1000<br>ELISA 1:10000   |
| <b>Formulation</b>         | Ascitic fluid, 0.03% Sodium Azide, 0.5% BSA, 50% Glycerol.   |
| <b>Isotype</b>             | IgG1   |
| <b>Storage Instruction</b> | Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.                         |

### TARGET INFORMATION

|                           |  |
|---------------------------|--|
| <b>Gene ID</b>            | 2268   |
| <b>Gene Symbol</b>        | FGR  |
| <b>Uniprot ID</b>         | FGR_HUMAN  |
| <b>Immunogen</b>          | Purified recombinant fragment of human c-Fgr expressed in E.coli.                                      |
| <b>Immunogen Region</b>   |  |
| <b>Specificity</b>        | FGR monoclonal antibody (Tyrosine-Protein Kinase Fgr) binds to endogenous Tyrosine-Protein Kinase Fgr. |
| <b>Immunogen Sequence</b> |  |



Immunohistochemistry analysis of paraffin-embedded human Spleen tissues with AEC staining using c-Fgr monoclonal antibody.



Western blot analysis using c-Fgr monoclonal antibody against HL60 (1) Raw264.7 (2) cell lysate.