

## Anti-NFATC2 antibody (640-700) (STJ98986)

STJ98986

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	Rabbit polyclonal antibody anti-Nuclear Factor Of Activated T-Cells-Cytoplasmic 2 (640-700) is suitable for use in Western Blot, Immunohistochemistry, Immunofluorescence and ELISA research applications.
<b>Applications</b>	WB, IHC-P, IF-P, ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human, Mouse, Rat

### PRODUCT PROPERTIES

<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	1 mg/mL
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	The antibody was affinity-purified from rabbit anti-serum by affinity-chromatography.
<b>Dilution</b>	IHC-P 1:50-200
<b>Range</b>	ELISA 1:10000-20000
<b>Formulation</b>	PBS, 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.
<b>Isotype</b>	IgG
<b>Storage</b>	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
<b>Instruction</b>	

### TARGET INFORMATION

<b>Gene ID</b>	<a href="#">4773</a>
<b>Gene Symbol</b>	<a href="#">NFATC2</a>
<b>Uniprot ID</b>	<a href="#">NFAC2_HUMAN</a>
<b>Immunogen</b>	Synthetic peptide from human protein at amino acid range: 640-700
<b>Immunogen Region</b>	640-700
<b>Specificity</b>	NFATC2 polyclonal antibody (Nuclear Factor Of Activated T-Cells-Cytoplasmic 2) binds to endogenous Nuclear Factor Of Activated T-Cells-Cytoplasmic 2 at the amino acid region 640-700.
<b>Immunogen Sequence</b>	



Immunohistochemical analysis of paraffin-embedded Human-colon, antibody was diluted at 1:100

---

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.  
St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081