

## Anti-OGT antibody (STJ99662)

STJ99662

### GENERAL INFORMATION


<b>Product Type</b>	Primary antibodies
<b>Short Description</b>	Rabbit polyclonal antibody anti-Udp-N-Acetylglucosamine--Peptide N-Acetylglucosaminyltransferase 110 Kda Subunit is suitable for use in Western Blot and ELISA research applications.
<b>Applications</b>	WB, ELISA
<b>Host/Source</b>	Rabbit
<b>Reactivity</b>	Human, Mouse, Rat

### PRODUCT PROPERTIES

<b>Clonality</b>	Polyclonal
<b>Clone ID</b>	
<b>Concentration</b>	1 mg/mL
<b>Conjugation</b>	Unconjugated
<b>Purification</b>	The antibody was affinity-purified from rabbit anti-serum by affinity-chromatography.
<b>Dilution</b>	WB 1:500-2000
<b>Range</b>	ELISA 1:10000-20000
<b>Formulation</b>	PBS, 50% Glycerol, 0.5% BSA and 0.02% Sodium Azide.
<b>Isotype</b>	IgG
<b>Storage</b>	Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
<b>Instruction</b>	

### TARGET INFORMATION

<b>Gene ID</b>	<a href="#">8473</a>
<b>Gene Symbol</b>	<a href="#">OGT</a>
<b>Uniprot ID</b>	<a href="#">OGT1_HUMAN</a>
<b>Immunogen</b>	Synthesized peptide derived from human OGT Polyclonal
<b>Immunogen Region</b>	
<b>Specificity</b>	OGT polyclonal antibody (Udp-N-Acetylglucosamine--Peptide N-Acetylglucosaminyltransferase 110 Kda Subunit) binds to endogenous Udp-N-Acetylglucosamine--Peptide N-Acetylglucosaminyltransferase 110 Kda Subunit.
<b>Immunogen Sequence</b>	



Western blot analysis of HEPG2 lysate, antibody was diluted at 1000. Secondary antibody was diluted at 1:20000