

Mouse Hepatocyte Growth Factor protein (Recombinant) (STJP000596)

STJP000596

GENERAL INFORMATION


Product Type Proteins
Short Description Recombinant-Mouse Hepatocyte Growth Factor-protein was developed from cho cells. For use in research applications.
Host/Source CHO cells

PRODUCT PROPERTIES


Concentration
Formulation Mouse HGF was lyophilized from a 0.2 μg/ml filtered solution in 2.5% glycine, 0.5% sucrose, 0.01% Tween80, 5 mM Glutamic acid, pH 4.5.
Purification
Dilution Range >97%, as determined by SDS-PAGE and HPLC
Storage Instruction The lyophilized protein is stable for at least 2 years from date of receipt at -20°C. Upon reconstitution, this cytokine can be stored in working aliquots at 2-8°C for one month, or at -20°C for six months, with a carrier protein without detectable loss

TARGET INFORMATION


Gene ID 15234
Gene Symbol Hgf
Uniprot ID HGF_MOUSE
Immunogen Sequence




BxPC-3 or ASPC-1 cells infected with recombinant lentivirus expressing Met knockdown shRNAs (2 or 5) or a non-targeting shRNA were treated without or with HGF and examined by Western analysis for pMet (Y1234/1235), Met, pErk1/2, Erk1/2, pAKT, AKT and Beta-actin levels (n=3).



Hepatic. Cells were seeded at a concentration of 5,000 cells/cm² on tissue culture plastic plates and coverslips coated with Matrigel and cultured in high glucose DMEM supplemented with 1% Penicillin/Streptomycin, 2 mmol/l L-Glutamine and 10% FBS for 3 days. The media were then changed to high glucose DMEM supplemented with 15% FBS, 1% Penicillin/Streptomycin, 2 mmol/l L-Glutamine, 300 μmol/l Monothioglycerol, 20 ng/ml Hepatocyte Growth Factor, 10 ng/ml Oncostatin M, 10⁻⁷ Dexamethasone, 100 ng/ml FGF4, and 1X ITS. The cells were allowed to differentiate for 21 days and then fixed and stored in PBS for immunofluorescence. The differentiation media were collected and analyzed for the presence of urea secreted by the differentiated cells. Urea was subsequently measured using the Urea/Ammonia determination kit (R-Biopharm AG, Darmstadt, Germany) according to the manufacturer's NA instructions. Cells were subsequently assessed for expression of the hepatocyte markers ALBUMIN and...



Prior to starting the hepatic differentiation, medium NA-MSCs at passage 5 were maintained in the regular medium NA-MSC culture medium until at 80-90% confluency. The hepatic differentiation was elicited in the differentiation-inducing medium, which consisted of MEM supplemented with 10% FBS, 10 ng/ml hepatocyte growth factor (HGF), 10 ng/ml basic fibroblast growth factor (bFGF) and 10 ng/ml oncostatin M (Osystem). The medium was changed every 3 days, and the cells were cultured for 8 days.



>Stage 3: Differentiation of Premature Hepatocyte-Like Cells (5-11) - At day 5, the medium was replaced by Knockout DMEM supplemented with 1% KSR, 1% nonessential amino acids, 1% l-glutamic acid, 1% dimethyl sulfoxide (DMSO), and 100 ng/ml hepatocyte growth factor (HGF) to induce premature hepatocyte-like cells. The medium was replaced every two days.

This product is suitable for in-vitro studies under the RESEARCH USE ONLY [RUO] licence. This product must not be used as for diagnostic or other medical purposes.
 St John's Laboratory Ltd, Knowledge Dock Business Centre, University Way, London, E16 2RD | Tel: 0208 223 3081