

Anti-Cd2ap antibody (360-639) (STJ110098)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 71kDa Observed Mw: 90kDa
Storage Instruction	Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	23607
Gene Symbol	CD2AP
UniProt ID	CD2AP_HUMAN
Immunogen Region	360-639
Immunogen Sequence	KYFSLKPEEKDEKSTLEQKP SKPAAPQVPPKKPTPTKAS NLLRSSGTVYPKRPEKPVPP PPPIAKINGEVSSISSKFET EPVSKLKLDSEQLPLRPKSV DFDSLTVRTSKETDVPNFD IASSENLLHLTANRPKMPGR RLPGRFNGGHSPTHSPEKIL KLPKEEDSANLKPSELKDDT CYSPKPSVYLSTPSSASKAN TTAFLTLEIKAKVETDDVK KNSLDELRAQIIELLCIVE
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 360-639 of human CD2AP (NP_036252.1).

ADDITIONAL INFORMATION

Note	STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.
------	---



Western blot analysis of various lysates using CD2AP Rabbit polyclonal antibody (STJ110098) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 60s.