

Anti-C8B antibody (55-350) (STJ110622)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 67kDa Observed Mw: 67kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	732
Gene Symbol	C8B
UniProt ID	C08B_HUMAN
Immunogen Region	55-350
Immunogen Sequence	SVDVTLMPIDCELSWSSWT TCDPCQKKRYRYAYLLQPSQ FHGEPNCFSDKEVEDCVNTR PCRSQVRCEGFVCAQTGRV NRRLCNGDNDCGDSDEAN CRRIYKCKQHEMDQYWGIGS LASGINLFTNSFEGPVLDR YYAGGCSPHYLNTRFRKPY NVESYTPQTQGYEFILKEY ESYSDFERNVTEKMASKSGF SFGFKIPGIFELGISSQSDR GKHYIRRTKRFSHTKSVFL
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 55-350 of human C8B (NP_000057.2).

ADDITIONAL INFORMATION

Note	STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.
------	---



Western blot analysis of various lysates using C8B Rabbit pAb (STJ110622) at 1:1000 dilution.
 Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution.
 Lysates/proteins: 25 µg per lane.
 Blocking buffer: 3% nonfat dry milk in TBST.
 Detection: ECL Basic Kit
 Exposure time: 90s.