

## Anti-EEA1 antibody (1182-1411) (STJ110997)

### GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

### PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 162kDa Observed Mw: 175kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

### TARGET INFORMATION


Gene ID	<a href="#">8411</a>
Gene Symbol	<a href="#">EEA1</a>
UniProt ID	<a href="#">EEA1_HUMAN</a>
Immunogen Region	1182-1411
Immunogen Sequence	LKAAVEQEKRNOQILKDQVK KEEELKKEFIEKEAKLHSE IKEKEVGMKKHEENEAKLTM QITALNENLGTVKKEWQSSQ RRVSELEKQTDLLRGEI AVL EATVQNNQDERRALLERCLK GEGEIEKLQTKVLELQRKLD NTTAAVQELGRENQSLQIKH TQALNRKWAEDNEVQNCMAC GKGFSVTVRRHHCRQCGNIF CAECSAKNALTPSSKKPVRV CDACFNDLQG
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1182-1411 of human EEA1 (NP_003557.2).

### ADDITIONAL INFORMATION


Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.




Immunohistochemistry analysis of paraffin-embedded Mouse kidney using EEA1 antibody (STJ110997) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded Human mammary cancer using EEA1 antibody (STJ110997) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded Rat lung using EEA1 antibody (STJ110997) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of various cell lines, using EEA1 antibody (STJ110997) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 5s.