

Anti-HILPDA antibody (1-63) (STJ11100178)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	IHC-P/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 7kDa Observed Mw: Refer to figures
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	29923
Gene Symbol	HILPDA
UniProt ID	HILPDA_HUMAN
Immunogen Region	1-63
Immunogen Sequence	MKHLNLYLLGVVLTLLSIF VRMESLEGLLESPSGTSW TTRSQLANTEPTKGLPDHPS RSM
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-63 of human HILPDA (NP_037464.1).

ADDITIONAL INFORMATION


Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



Immunohistochemistry analysis of HILPDA in paraffin-embedded Mouse liver using HILPDA Rabbit polyclonal antibody (STJ11100178) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of HILPDA in paraffin-embedded Human colon using HILPDA Rabbit polyclonal antibody (STJ11100178) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of HILPDA in paraffin-embedded Rat liver using HILPDA Rabbit polyclonal antibody (STJ11100178) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.