

## Anti-PSMD6 antibody (23-307) (STJ11100219)

### GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

### PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 46kDa Observed Mw: 46kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

### TARGET INFORMATION


Gene ID	<a href="#">9861</a>
Gene Symbol	<a href="#">PSMD6</a>
UniProt ID	<a href="#">PSMD6_HUMAN</a>
Immunogen Region	23-307
Immunogen Sequence	RFLLSLPEHRGDAAVRDELMAAVRDNNMAPYYEALCKSLD WQIDVDLLNKMKKANEDELK RLDEELEDKAEKNGESEIRD AMMAKAEYLCRIGDKEGALT AFRKTYDKTVALGHRLDIVF YLLRIGLFYMDNDLITRNTKAKSLIEEGDWDRRRLKLV YQGLYCVAIRDFKQAAELFL DTVSTFTSYELMDYKTFVTY TVYVSMIALERPDLREKVIK GAEILEVLHSLPAVRQYLF
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 23-307 of human PSMD6 (NP_055629.1).

### ADDITIONAL INFORMATION

Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



Western blot analysis of lysates from Rat brain, using PSMD6 Rabbit polyclonal antibody (STJ11100219) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 90s.



Western blot analysis of various lysates using PSMD6 Rabbit polyclonal antibody (STJ11100219) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 3min.