

Anti-SIRT6 antibody (1-355) (STJ11100422)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 39kDa Observed Mw: 39kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	51548
Gene Symbol	SIRT6
UniProt ID	SIR6_HUMAN
Immunogen Region	1-355
Immunogen Sequence	MSVNYAAGLSPYADKGGKGL PEIFDPPEELERKVVWELARL VWQSSVVFHTGAGISTASG IPDFRGPVHVTMEERGLAP KFDTTFESARPTQTHMALVQ LERVGLLRFLVSNVDGLHV RSGFPRDKLAEHLHGNMFVEE CAKCKTQYVRDVTVGTMLGK ATGRLCTVAKARGLRACRGE LRDTILDWEDSLPDRDLALA DEASRNADLSITLGTSLQIR PSGNPLATKRRGGRLVIV
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-355 of human SIRT6 (NP_057623.2).

ADDITIONAL INFORMATION

Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma using SIRT6 Rabbit polyclonal antibody (STJ11100422) at dilution of 1:50 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of various cell lines, using SIRT6 antibody (STJ11100422) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.