



Anti-Tri-Methyl-Histone H3-K64 antibody (STJ11100934)


GENERAL INFORMATION	
Product Type	Primary antibodies
Applications	WB/IHC-P/IF/ICC/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat/Other
PRODUCT PROPERTIES	
Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IHC-P:1:50-1:200 IF/CC:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 16kDa Observed Mw: 16kDa/17kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.
TARGET INFORMATION	
Gene ID	8350/8351/8352/8353/8354/8355/8356/8357/8358/8359/8360
Gene Symbol	H3C1.H3C2.H3C3.H3C4.H3C6.H3C7.H3C8.H3C10.H3C11.H3C12 H31_HUMAN H31T_HUMAN
Immunogen Sequence	YQKST
Specificity	A synthetic trimethylated peptide around K56 of human TriMethyl-Histone H3-K64 (NP_003520.1).
ADDITIONAL INFORMATION	
Note	STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.




Immunofluorescence analysis of NIH/3T3 cells using TriMethyl-Histone H3-K64 Rabbit polyclonal antibody (STJ11100934) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of HeLa cells using TriMethyl-Histone H3-K64 Rabbit polyclonal antibody (STJ11100934) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of TriMethyl-Histone H3-K64 in paraffin-embedded mouse brain using TriMethyl-Histone H3-K64 Rabbit polyclonal antibody (STJ11100934) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.




Immunohistochemistry analysis of TriMethyl-Histone H3-K64 in paraffin-embedded human breast using TriMethyl-Histone H3-K64 Rabbit polyclonal antibody (STJ11100934) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of TriMethyl-Histone H3-K64 in paraffin-embedded rat brain using TriMethyl-Histone H3-K64 Rabbit polyclonal antibody (STJ11100934) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of various lysates using TriMethyl-Histone H3-K64 Rabbit polyclonal antibody (STJ11100934) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 120s.



Western blot analysis of various lysates using TriMethyl-Histone H3-K64 Rabbit polyclonal antibody (STJ11100934) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 150s.