

Anti-VDAC2 antibody (14-24) (STJ11101478)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/IF/ICC/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IHC-P:1:50-1:200 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 32kDa Observed Mw: 35kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	7417
Gene Symbol	VDAC2
UniProt ID	VDAC2_HUMAN
Immunogen Region	14-24
Immunogen Sequence	MATHGQTCARPMCIPPSYAD LGKAARDIFNKGFGLVVL DVKTKSCSGVEFSTSGSSNT DTGKVTGTLETQYKWCEYGL TFTEKWNTDNTLGTETIAIED
Specificity	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human VDAC2 (NP_003366.2).

ADDITIONAL INFORMATION


Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.




Immunofluorescence analysis of PC-12 cells using VDAC2 Rabbit polyclonal antibody (STJ11101478) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




Immunofluorescence analysis of NIH/3T3 cells using VDAC2 Rabbit polyclonal antibody (STJ11101478) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of paraffin-embedded rat heart using VDAC2 Rabbit polyclonal antibody (STJ11101478) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.




Immunohistochemistry analysis of paraffin-embedded mouse testis using VDAC2 Rabbit polyclonal antibody (STJ11101478) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded human breast cancer using VDAC2 Rabbit polyclonal antibody (STJ11101478) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded human brain using VDAC2 Rabbit polyclonal antibody (STJ11101478) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of various cell lines, using VDAC2 antibody (STJ11101478) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.