

Anti-RAP1GAP antibody (1-100) [S2MR] (STJ11101912)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IF/ICC/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Monoclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IF/CC:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 73kDa Observed Mw: 95kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	5909
Gene Symbol	RAP1GAP
UniProt ID	RPGP1_HUMAN
Immunogen Region	1-100
Immunogen Sequence	MIEKMQGSRMDEQRCSEFPPP LKTEEDYIPYPSVHEVLGRE GPFPLILLPQFGGYWIEGTN HEITSIPETEPLQSPPTTKVK LECNPTARIYRKHFGLGKEHF
Specificity	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Rap1GAP (P47736).

ADDITIONAL INFORMATION

Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



Immunofluorescence analysis of NIH-3T3 cells using Rap1GAP Rabbit monoclonal antibody (STJ11101912) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.



Western blot analysis of extracts of various cell lines, using Rap1GAP Rabbit monoclonal antibody (STJ11101912) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 3s.