

Anti-MYLK antibody (1800-1900) [S7MR] (STJ11101957)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/ELISA
Host / Source	Rabbit
Reactivity	Human/Rat

PRODUCT PROPERTIES


Clonality	Monoclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 ELISA: Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 211kDa Observed Mw: 130kDa/140kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	4638
Gene Symbol	MYLK
UniProt ID	MYLK_HUMAN
Immunogen Region	1800-1900
Immunogen Sequence	VAEEKPHVKPYFSKTIRDLE VVEGSAARFDCKIEGYDPPE VVWFKDDQSIRESRHFQIDY DEDGNCSLIISDVC GDDDAK YTCKAVNSLGEATCTAELIV E
Specificity	A synthetic peptide corresponding to a sequence within amino acids 1800-1900 of human MYLK (Q15746).

ADDITIONAL INFORMATION

Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.



Western blot analysis of extracts of various cell lines, using MYLK Rabbit monoclonal antibody (STJ11101957) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.