

Anti-ADORA1 antibody (100-200) [S9MR] (STJ11102299)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Monoclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:1000-1:6000 IHC-P:1:100-1:1000 ELISA:Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 37kDa Observed Mw: 35kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	134
Gene Symbol	ADORA1
UniProt ID	AA1R_HUMAN
Immunogen Region	100-200
Immunogen Sequence	AIAVDRLRVKIPRLRYKMVV TPRRAAVAAGCWILSFVVG LTPMFGWNNLSAVERAWAAN GSMGEPVIKCEFEKVISM EY M VYFNFFVWVLP LLLMVL I Y
Specificity	A synthetic peptide corresponding to a sequence within amino acids 100-200 of human ADORA1 (P30542).

ADDITIONAL INFORMATION


Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**




Immunohistochemistry analysis of ADORA1 in paraffin-embedded mouse spinal cord using ADORA1 Rabbit monoclonal antibody (STJ11102299) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of ADORA1 in paraffin-embedded human brain using ADORA1 Rabbit monoclonal antibody (STJ11102299) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of ADORA1 in paraffin-embedded rat brain using ADORA1 Rabbit monoclonal antibody (STJ11102299) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with immunohistochemistry staining protocol.



Western blot analysis of various lysates using ADORA1 Rabbit monoclonal antibody (STJ11102299) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.