

Anti-NDST1 antibody (40-170) (STJ11102911)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 101kDa Observed Mw: 101kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	3340
Gene Symbol	NDST1
UniProt ID	NDST1_HUMAN
Immunogen Region	40-170
Immunogen Sequence	GWKRGLEPSADAPEPCGDP PPVAPSRLPLKPVQAATPS RTDPLVLFVESLYSQLGQE VVAILESSRFKYRTEIAPGK GDMPTLTDKGRGRFALIYE NILKYVNLDAWNRELLDKYC VAYGVGIIGFF
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 40-170 of human NDST1 (NP_001534.1).

ADDITIONAL INFORMATION

Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.



Western blot analysis of lysates from HepG2 cells, using NDST1 Rabbit polyclonal antibody (STJ11102911) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.