

Anti-DNAJC17 antibody (1-304) (STJ11103199)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 35kDa Observed Mw: 35kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	55192
Gene Symbol	DNAJC17
UniProt ID	DJC17_HUMAN
Immunogen Region	1-304
Immunogen Sequence	MAVTKELLQMDLYALLGIEE KAADKEVKKAYRQKALSCHP DKNPDPNPRAAELFHQLSQAL EVLTDAAARAAYDKVRKAKK QAAERTQKLDKRRKKVLDL EARERQAQAQEEEEESRS TRTLEQEIERLREEGSRQLE EQQLIREQIRQERDQRLRG KAENTEGGGTPKLLKWKCK KEDESKGGYSKDVLLRLLQK YGEVNLNLVLSKKPGTAVVE FATVKAAELAVQNEVGLVD
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-304 of human DNAJC17 (NP_060633.1).

ADDITIONAL INFORMATION


Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.




Immunohistochemistry analysis of DNAJC17 in paraffin-embedded mouse spinal cord using DNAJC17 Rabbit polyclonal antibody (STJ11103199) at dilution of 1:20 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of DNAJC17 in paraffin-embedded human esophageal cancer using DNAJC17 Rabbit polyclonal antibody (STJ11103199) at dilution of 1:20 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Western blot analysis of various lysates using DNAJC17 Rabbit polyclonal antibody (STJ11103199) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 180s.



Western blot analysis of various lysates using DNAJC17 Rabbit polyclonal antibody (STJ11103199) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 180s.