# Anti-Periostin antibody (1-100) [S2MR] (STJ11103642)

## GENERAL INFORMATION

| | |
|---|---|
| Product Type | Primary antibodies |
| Applications | WB/ELISA |
| Host / Source | Rabbit |
| Reactivity | Human |

## PRODUCT PROPERTIES


| | |
|---|---|
| Clonality | Monoclonal |
| Concentration | Lot specific |
| Conjugation | Unconjugated |
| Purification | Affinity purification |
| Dilution Range | WB:1:500-1:1000<br>ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements. |
| Formulation | PBS with 0.02% Sodium Azide, 0.05% BSA, 50% Glycerol, pH 7.3. |
| Isotype | IgG |
| Molecular Weight | Protein Mw: 80kDa/83kDa/84kDa/87kDa/90kDa/93kDa<br>Observed Mw: 80kDa |
| Storage Instruction | Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles. |

## TARGET INFORMATION

| | |
|---|---|
| Gene ID | 10631 |
| Gene Symbol | POSTN |
| UniProt ID | POSTN_HUMAN |
| Immunogen Region | 1-100 |
| Immunogen Sequence | MIPFLPMFSLLLLLIVNPIN ANNHYDKILAHSRIRGRDQG PNVCALQQILGTKKKYFSTC KNWYKKSICGQKTTVLYECC PGYMRMEGMKGCPAVLPIDH |
| Specificity | A synthetic peptide corresponding to a sequence within amino acids 1-100 of human Periostin (NP_001129406.1). |

## ADDITIONAL INFORMATION

| | |
|---|---|
| Note | STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS. |




Western blot analysis of lysates from MCF7 cells, using Periostin Rabbit mAb (STJ11103642) at 1:1000 dilution.
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution.
Lysates/proteins: 25 Mu g per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit
Exposure time: 1s.