

Anti-PIK3C2A antibody (1-200) (STJ11104405)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/IF/ICC/IP/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IHC-P:1:50-1:200 IF/CC:1:50-1:200 IP:0.5 Mu g-4 Mu g antibody for 200 Mu g-400 Mu g extracts of whole cells ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specif
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 191kDa Observed Mw: Refer to figures
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	5286
Gene Symbol	PIK3C2A
UniProt ID	P3C2A_HUMAN
Immunogen Region	1-200
Immunogen Sequence	MAQISSNSGFKCEKPPSSHPEP TRAKDVKKEALQMEAEALA KLQKDRQVTDNQRGFEFLSSS TRKKAQVYNKQDYDLMVFPE SDSQKRALDIDVEKLTQAEEL EKLLDDSFETKKTPLVLPVT PILSPSFSQAQLYFRPTIQRG QWPPLGPGPSTYALPSIYPS TYSKQAFAFQNGFNPRMPTFP STEPIYLSLPGQSPYFYSYPL
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human PIK3C2A (NP_002636.2).

ADDITIONAL INFORMATION


Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.




Immunoprecipitation analysis of 200 Mu g extracts of HeLa cells, using 3 Mu g PIK3C2A antibody (STJ11104405). Western blot was performed from the immunoprecipitate using PIK3C2A antibody (STJ11104405) at a dilution of 1:1000.




Immunofluorescence analysis of C6 cells using PIK3C2A Rabbit polyclonal antibody (STJ11104405) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of PIK3C2A in paraffin-embedded human liver damage using PIK3C2A Rabbit polyclonal antibody (STJ11104405) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of PIK3C2A in paraffin-embedded rat heart using PIK3C2A Rabbit polyclonal antibody (STJ11104405) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of PIK3C2A in paraffin-embedded mouse heart using PIK3C2A Rabbit polyclonal antibody (STJ11104405) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of various lysates using PIK3C2A Rabbit polyclonal antibody (STJ11104405) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.