

Anti-ECD antibody (435-644) (STJ11104764)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 73kDa Observed Mw: 76kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	11319
Gene Symbol	ECD
UniProt ID	ECD_HUMAN
Immunogen Region	435-644
Immunogen Sequence	ESESVSKEEKEQNYDLTEVS ESMKAFISKVSTHKGAE LPR EPSEAPITFDADSFLNYFDK ILGPRPNESDSDLDDEDFE CLDSDDDLDFETHEPGEEAS LKGTLDNLKSYMAQMDQELA HTCISKSFTRNQVEPVSTQTDNNSDEEDSGTGESVMPV DVDLNLVSNILESYSSQAGL AGPASNLLQSMGVQLPDNTD HRPTSKPTKN
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 435-644 of human ECD (NP_009196.1).

ADDITIONAL INFORMATION

Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



Western blot analysis of various lysates, using ECD antibody (STJ11104764) at 1:800 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.



Western blot analysis of HeLa, using ECD antibody (STJ11104764) at 1:800 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.