

## Anti-MKP1 antibody (1-172 aa) (STJ11105073)

### GENERAL INFORMATION

|               |                    |
|---------------|--------------------|
| Product Type  | Primary antibodies |
| Applications  | WB/ELISA           |
| Host / Source | Rabbit             |
| Reactivity    | Human/Mouse/Rat    |

### PRODUCT PROPERTIES



|                     |   |
|---------------------|---|
| Clonality           | Polyclonal  |
| Concentration       | Lot specific  |
| Conjugation         | Unconjugated  |
| Purification        | Affinity purification   |
| Dilution Range      | WB:1:1000-1:5000<br>ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements. |
| Formulation         | PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.  |
| Isotype             | IgG   |
| Molecular Weight    | Protein Mw: 39kDa<br>Observed Mw: 40kDa   |
| Storage Instruction | Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.  |

### TARGET INFORMATION

|                    |   |
|--------------------|---|
| Gene ID            | <a href="#">19252</a>   |
| Gene Symbol        | <a href="#">Dusp1</a>   |
| UniProt ID         | <a href="#">DUS1_MOUSE</a>  |
| Immunogen Region   | 1-172 aa  |
| Immunogen Sequence | MVMEVGILDAGGLRALLREG AAQCLLLDCRSFFAFNAGHI AGSVNVRVFSTIVRRRAKGAM<br>GLEHIVPNAELRGRLLAGAY HAVVLLDERSASLDGAKRDG TLALAAGALCREARSTQVFF<br>LQGGYEAFSASCPELCSKQS TPTGLSLPLSTVPSAESG CSSCSTPLYDQG |
| Specificity        | Recombinant fusion protein containing a sequence corresponding to amino acids 1-172 of mouse MKP1. (NP_038670.1).   |

### ADDITIONAL INFORMATION

Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.



Western blot analysis of various lysates, using MKP1 Rabbit polyclonal antibody (STJ11105073) at 1:2000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 2s.