

Anti-BRD9 antibody (274-597) (STJ11105726)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 67kDa Observed Mw: 80kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	65980
Gene Symbol	BRD9
UniProt ID	BRD9_HUMAN
Immunogen Region	274-597
Immunogen Sequence	REVISCMFEPEGNACSLTDS TAEHVLALVEHADEARDR INRFLPGGKMGYLKRNGDGS LLYSVNTAEPDADEEETHP VDLSSLSSKLLPGFTTLGFK DERRNKVTLSSATTALSMQ NNSVFGDLKSDMELLYSAY GDETVQCALSLOEFVKDAG SYSKKVVDLLDQITGGDHS RTLFLQLKQRRNVPMPKPDEA KVGDTLGDSSSSVLEFMSMK SYPDVSVDISMLSSLGKVK
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 274-597 of human BRD9 (NP_076413.3).

ADDITIONAL INFORMATION


Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.



Immunohistochemistry analysis of paraffin-embedded mouse brain using BRD9 antibody (STJ11105726) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded human colon using BRD9 antibody (STJ11105726) at dilution of 1:200 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of various cell lines, using BRD9 antibody (STJ11105726) at 1:500 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 3s.