

Anti-HA-Tag antibody [SMM5506] {HRP} (STJ11107576)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/ELISA
Host / Source	Mouse
Reactivity	Species Independent

PRODUCT PROPERTIES


Clonality	Monoclonal
Concentration	Lot specific
Conjugation	HRP
Purification	Affinity purification
Dilution Range	WB:1:500-1:5000 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 50% Glycerol, pH 7.3.
Isotype	IgG1k
Molecular Weight	53kDa (CD95) /65kDa (CD2) /53kDa (SCRN1) /26kD (HA (N端)) /42kDa (WDR77) /45kDa (NR3C2)

TARGET INFORMATION


Immunogen Sequence	YPYDVPDYA
Specificity	A synthetic peptide corresponding to HA tag.

ADDITIONAL INFORMATION


Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



Western blot analysis of extracts of normal 293T cells, 293T transfected with SCR1 Protein, using HRP-conjugated Mouse anti HA-Tag mAb (STJ11107576) at 1:5000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.



Western blot analysis of extracts of normal 293T and 293F cells, 293T transfected with CD2 Protein and HA-N Protein, 293F transfected with CD95-N Protein using HRP-conjugated Mouse anti HA-Tag mAb (STJ11107576) at 1:5000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 3s.



Western blot analysis of extracts of normal 293T and 293F cells, 293T transfected with NR3C2-C Protein, 293F transfected with WDR77-C Protein using HRP-conjugated Mouse anti HA-Tag mAb (STJ11107576) at 1:5000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.