

## Anti-DLG1 antibody (1-190) (STJ111281)

### GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IF/ICC/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

### PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IF/CC:1:50-1:200 ELISA:Recommended starting concentration is 1 $\mu$ g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 100kDa Observed Mw: 114kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

### TARGET INFORMATION


Gene ID	<a href="#">1739</a>
Gene Symbol	<a href="#">DLG1</a>
UniProt ID	<a href="#">DLG1_HUMAN</a>
Immunogen Region	1-190
Immunogen Sequence	MPVVRKQDTQRALHLLSEYRS KLSQTEDRQLRSSIERVINI FQSNLFQALIDIQEFYEVTL LDNPKCIDRSKPSEPIQPVN TWEISSLPSTVTSETLPSS LSPSVEKYRYQDEDETPPQEH ISPQITNEVIGPELVHVHSEK NLSEIENVHGFVSHSHISPI KPTEAVLPSPPTVPVIVLP VPAENTVILP
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-190 of human DLG1 (NP_004078.2).

### ADDITIONAL INFORMATION

Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



Western blot analysis of various lysates using DLG1 Rabbit pAb (STJ111281) at 1:1000 dilution. Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution.  
Lysates/proteins: 25  $\mu$ g per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit  
Exposure time: 90s.



Immunofluorescence analysis of L929 cells using DLG1 Rabbit pAb (STJ111281) at dilution of 1:100 (40x lens). Secondary antibody: Cy3-conjugated Goat anti-Rabbit IgG (H+L) (STJS001166) at 1:500 dilution. Blue: DAPI for nuclear staining.