

## Anti-CCNK antibody (1-160) (STJ112299)

### GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

### PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 64kDa Observed Mw: 90kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

### TARGET INFORMATION


Gene ID	<a href="#">8812</a>
Gene Symbol	<a href="#">CCNK</a>
UniProt ID	<a href="#">CCNK_HUMAN</a>
Immunogen Region	1-160
Immunogen Sequence	MKENKENSPPSVTSANLDHT KPCWYWDKDLAHTPSQLEG LDPATEARYRREGARFIDV GTRLGLHYDTLATGIIYFHR FYMFHSFKQFPRYVTGACCL FLAGKVEETPKKCKDIIKTA RSLNDVQFGQFGDDPKKEV MVLERILLQTIKFDLQVEHP
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-160 of human Cyclin K (NP_001092872.1).

### ADDITIONAL INFORMATION


Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**




Immunohistochemistry analysis of paraffin-embedded rat brain using Cyclin K Rabbit polyclonal antibody (STJ112299) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded mouse spinal cord using Cyclin K Rabbit polyclonal antibody (STJ112299) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded human breast cancer using Cyclin K Rabbit polyclonal antibody (STJ112299) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of various cell lines, using Cyclin K Antibody (STJ112299) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 90s.