

Anti-NCOA2 antibody (550-820) (STJ112318)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/ELISA
Host / Source	Rabbit
Reactivity	Human/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 159kDa Observed Mw: 200kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	10499
Gene Symbol	NCOA2
UniProt ID	NCOA2_HUMAN
Immunogen Region	550-820
Immunogen Sequence	SSLASPDCLKMGNLQNSPVNM NPPPLSKMGSLDSKDCFGLY GEPSEGTGQAESSCHPGEQ KETNDPNLPPAVSSERADGQ SRLHDSKGGTKLLQLLTTKS DQMEPSPPLASSLSDTNKDST GSLPGSGSTHGTSLKEKHKI LHRLLQDSSSPVDLAKLTAE ATGKDLQSQSSSTAPGSEVT IKQEPVSPKKKENALLRYLL DKDDTKDIGLPEITPKLERL DSKTDPASNTKLIAMKTEK
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 550-820 of human NCOA2 (NP_006531.1).

ADDITIONAL INFORMATION


Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**




Immunohistochemistry analysis of paraffin-embedded human esophageal cancer using [KO Validated] NCOA2 Rabbit polyclonal antibody (STJ112318) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat lung using [KO Validated] NCOA2 Rabbit polyclonal antibody (STJ112318) at dilution of 1:100 (40x lens). Perform high pressure antigen retrieval with 10 mM citrate buffer pH 6.0 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of various cell lines, using NCOA2 antibody (STJ112318) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 180s.



Western blot analysis of extracts from normal (control) and NCOA2 knockout (KO) 293T cells, using NCOA2 antibody (STJ112318) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 3min.