Anti-NACC1 antibody (170-340) (STJ113847)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 57kDa Observed Mw: 57kDa
Storage Instruction	Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	112939
Gene Symbol	NACC1
UniProt ID	NACC1_HUMAN
Immunogen Region	170-340
Immunogen Sequence	QQESDSVQCMPVAKRLWDSG QKEAGGGGNGSRKMAKFSTP DLAANRPHQPPPPQQAPVVA AAQPAVAAGAGQPAGGVAAA GGVVSGPSTSERTSPGTSSA YTSDSPGSYHNEEDEEEDGG EEGMDEQYRQICNMYTMYSM MNVGQTAEKVEALPEQVAPE SRNRIRVRQDL
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 170-340 of human NACC1 (NP_443108.1).

ADDITIONAL INFORMATION

Note	STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.



Western blot analysis of various lysates using NACC1 Rabbit polyclonal antibody (STJ113847) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 10s.