

Anti-HNRNPL antibody (282-589) (STJ114321)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/IF/ICC/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IHC-P:1:50-1:200 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 64kDa Observed Mw: 64kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	3191
Gene Symbol	HNRNPL
UniProt ID	HNRPL_HUMAN
Immunogen Region	282-589
Immunogen Sequence	TWDTYTNPNLSGGQDPSNPN KRQRQPPLLDHPAEYGGPH GGYHSHYHDEGYGPPPPHYE GRRMGPPVGGHRRGSPRYGP QYGHPPPPPPPEYGP HADS PVLMMVYGLDQSKMNCDRVFN VFCLYGNVEKVKFMKSKPGA AMVEMADGYAVDRAITHLNN NFMFGQKLNVCVSKQPAIMP GQSYGLEDGSCSYKDFSESR NNRFSTPEQAAKNRIQHPSN VLHFFNAPLEVTEENFFEI
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 282-589 of human HNRNPL (NP_001524.2).

ADDITIONAL INFORMATION


Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.




Immunofluorescence analysis of 293T cells using HNRNPL antibody (STJ114321) at dilution of 1:100 (40x lens). Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of paraffin-embedded mouse brain using HNRNPL antibody (STJ114321) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded human placenta using HNRNPL antibody (STJ114321) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of paraffin-embedded rat lung using HNRNPL antibody (STJ114321) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7. 2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of extracts of various cell lines, using HNRNPL antibody (STJ114321) at 1:10000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.