

Anti-GARS antibody (60-370) (STJ115269)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IF/ICC/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IF/CC:1:50-1:200 ELISA:Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.02% Sodium Azide, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 83kDa Observed Mw: 70-83kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	2617
Gene Symbol	GARS1
UniProt ID	GARS_HUMAN
Immunogen Region	60-370
Immunogen Sequence	AEEVLAPLRLAVRQQGDLVR KLKEDKAPQVDVDKAVAEK ARKRVLEAKELALQPKDDIV DRAKMEDTLKRRFFYDQAF A IYGGVSGLYDFGVPVGCALKN NIIQTWRQHFQEEQILEID CTMLTPEPVLKTSGHVDKFA DFMVKDVKNGECFRADHLLK AHLQKLMSDKKCSVEKKSEM ESVLAQLDNYGQQELADLFV NYNVKSPITGNDLSPVVSFN LMFKTFIGPGGNMPGYLRP
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 60-370 of human GARS (NP_002038.2).

ADDITIONAL INFORMATION

Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



Western blot analysis of extracts of various cell lines, using GARS antibody (STJ115269) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 5s.