

Anti-CD68 antibody (16-200) (STJ117231)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IF/ICC/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IF/CC:1:50-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 37kDa Observed Mw: 70-80kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	968
Gene Symbol	CD68
UniProt ID	CD68_HUMAN
Immunogen Region	16-200
Immunogen Sequence	AAQGTGNDPCPHKKSATLLPS FTVTPTVTESTGTTSHRTTK SHKTTTTHRTTTTGTTSHGPT TATHNPPTTSHGNVTVHPTS NSTATSQGPSTATHSPATTS HGNATVHPTSNSTATSPGFT SSAHPEPPPPSPSPSPSPTSKE TIGDYTWNGSQPCVHLQAQ IQIRVMYTTQGGGEAWGISV LNPKN
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 16-200 of human CD68 (NP_001242.2).

ADDITIONAL INFORMATION

Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



Immunofluorescence analysis of THP-1 cells using CD68 Rabbit polyclonal antibody (STJ117231) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using CD68 Rabbit polyclonal antibody (STJ117231) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.