

Anti-RPAP3 antibody (17-241) (STJ117433)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IF/ICC/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 76kDa Observed Mw: 76kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	79657
Gene Symbol	RPAP3
UniProt ID	RPAP3_HUMAN
Immunogen Region	17-241
Immunogen Sequence	NAEELQDFMRDLENWEKDIK QKDMELRRQNGVPEENLPPI RGNFRKKKKKGKAKESSKKT REENTKNRIKSYDYEAQWL DVDRILDELDDKDDSTHESLS QESESEEDGIHVDSQKALVL KEKGNKYFKQGGYDEAIDCY TKGMDADPYNPVLPNTNRASA YFRLKKFAVAESDCNLAVAL NRSYTKAYSRRGAARFALQK LEEAKDYEVRVLELEPNFE ATNEL
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 17-241 of human RPAP3 (NP_001139547).

ADDITIONAL INFORMATION


Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.



Immunofluorescence analysis of L929 cells using RPAP3 Rabbit polyclonal antibody (STJ117433) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of HeLa cells using RPAP3 Rabbit polyclonal antibody (STJ117433) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using RPAP3 Rabbit polyclonal antibody (STJ117433) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 1s.