

Anti-PLS1 antibody (1-110) (STJ117498)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:200-1:2000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 70kDa Observed Mw: 70kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	5357
Gene Symbol	PLS1
UniProt ID	PLS1_HUMAN
Immunogen Region	1-110
Immunogen Sequence	MENSTTTISREELEELQEAF NKIDIDNSGYVSDYELQDLF KEASLPLPGYKVVREIVEKIL SVADSNKDGKISFEFVSLM QELKSKDISKTFRKIINKRE GITAIGGTST
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-110 of human PLS1 (NP_001165783.1).

ADDITIONAL INFORMATION


Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.



Immunohistochemistry analysis of PLS1 in paraffin-embedded mouse kidney using PLS1 Rabbit polyclonal antibody (STJ117498) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of PLS1 in paraffin-embedded rat kidney using PLS1 Rabbit polyclonal antibody (STJ117498) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of various lysates using PLS1 Rabbit polyclonal antibody (STJ117498) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 3s.