

Anti-TPD52L3 antibody (1-132) (STJ117740)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IF/ICC/ELISA
Host / Source	Rabbit
Reactivity	Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:200-1:2000 IF/CC:1:50-1:200 ELISA:Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 16kDa Observed Mw: 14kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	89882
Gene Symbol	TPD52L3
UniProt ID	TPD55_HUMAN
Immunogen Region	1-132
Immunogen Sequence	MPHARTETSVGTYESHSTSE LEDLTEPEQRELKTKLTKLE AEIVTLRHVLAAKERRCGEL KRKLGTLTALVGLRQNLKSKSW LDVQVSNITYVKQKTSAAALST MGLTICRKLGGVKKSATLRS FEGNPKGEGSRI
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-132 of human TPD52L3 (NP_001001875.1).

ADDITIONAL INFORMATION


Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.



Immunofluorescence analysis of paraffin-embedded mouse testis using TPD52L3 Rabbit polyclonal antibody (STJ117740) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of paraffin-embedded rat testis using TPD52L3 Rabbit polyclonal antibody (STJ117740) at dilution of 1:100. Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.



Western blot analysis of various lysates using TPD52L3 Rabbit polyclonal antibody (STJ117740) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30S.