

## Anti-AQR antibody (400-700) (STJ117823)

### GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

### PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 IHC-P:1:50-1:100 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 171kDa Observed Mw: 171kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

### TARGET INFORMATION


Gene ID	<a href="#">9716</a>
Gene Symbol	<a href="#">AQR</a>
UniProt ID	<a href="#">AQR_HUMAN</a>
Immunogen Region	400-700
Immunogen Sequence	KEFLLELLVSRHERRISQIQQLNQMPLYPTEKIIWDENIVPTEYYSGEGCLALPKLNLQFLTLHDYLLRNFNLFRLSEYIEIQDIEDSVSRMKPWQSEYGGVVFGGWARMAPQIVAFTVVEVAKPNIGENWPTRVRADV TINLNVDRDHKDEWEGLRKH DVCFLITVRPTKPYGTFKFDRRPFIEQVGLVYVRGCEIQGMLDDKGRVIEDGPEPRPNLRGESRTFRVFLDPNQYQQDM
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 400-700 of human AQR (NP_055506.1).

### ADDITIONAL INFORMATION


Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.




Immunohistochemistry analysis of AQR in paraffin-embedded mouse testis using AQR Rabbit polyclonal antibody (STJ117823) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of AQR in paraffin-embedded human lung cancer using AQR Rabbit polyclonal antibody (STJ117823) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of AQR in paraffin-embedded rat spleen using AQR Rabbit polyclonal antibody (STJ117823) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of various lysates using AQR Rabbit polyclonal antibody (STJ117823) at 1:3000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 40s.