

## Anti-HACE1 antibody (1-100) (STJ117867)

### GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

### PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 102kDa Observed Mw: 102kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

### TARGET INFORMATION


Gene ID	<a href="#">57531</a>
Gene Symbol	<a href="#">HACE1</a>
UniProt ID	<a href="#">HACE1_HUMAN</a>
Immunogen Region	1-100
Immunogen Sequence	MERAMEQLNRLTRSLRRART VEPEDNETAVYTLMPMVMA DQHRSVSELLSNSKFDVNYA FGRVKRSLLLHIAANCGSVEC LVLLKKGANPNYQDISGCT
Specificity	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human HACE1 (NP_065822.2).

### ADDITIONAL INFORMATION


Note STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.



Western blot analysis of various lysates using HACE1 Rabbit pAb (STJ117867) at 1:1000 dilution.  
Secondary antibody: HRP-conjugated Goat anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution.  
Lysates/proteins: 25 µg per lane.  
Blocking buffer: 3% nonfat dry milk in TBST.  
Detection: ECL Basic Kit  
Exposure time: 10s.



Immunohistochemistry analysis of paraffin-embedded Human placenta using HACE1 Rabbit pAb (STJ117867) at dilution of 1:100 (40x lens).  
Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to immunohistochemistry staining.



Immunohistochemistry analysis of paraffin-embedded Rat testis using HACE1 Rabbit pAb (STJ117867) at dilution of 1:100 (40x lens).  
Microwave antigen retrieval performed with 0.01M PBS Buffer (pH 7.2) prior to immunohistochemistry staining.