

Anti-TRIM17 antibody (280-477) (STJ118281)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:1000 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.05% Proclin300, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 54kDa Observed Mw: 54kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION

Gene ID	51127
Gene Symbol	TRIM17
UniProt ID	TRI17_HUMAN
Immunogen Region	280-477
Immunogen Sequence	VCRVPGQIEVLRGFLEDVVP DATSAYPYLLLYESRQRRYL GSSPEGSGFCSKDRFVAYPC AVGQTAFSSGRHYWEVGMNI TGDALWALGVCRDNVSRKDR VPKCPENGFVVVQLSKGTKY LSTFSALTPVMLMEPPSHMG IFLDFEAGEVSFYVSVDGSH LHTYSQATFPGLQPFVCLG APKSGQMIVISTVMWVKG
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 280-477 of human TRIM17 (NP_057186.1).

ADDITIONAL INFORMATION

Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



Western blot analysis of various lysates, using TRIM17 antibody (STJ118281) at 1:400 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 µg per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 30s.