

Anti-TRIM27 antibody (1-200) (STJ118505)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	IHC-P/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	IHC-P:1:50-1:200 ELISA:Recommended starting concentration is 1 µg/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 58kDa Observed Mw: 58kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	5987
Gene Symbol	TRIM27
UniProt ID	TRI27_HUMAN
Immunogen Region	1-200
Immunogen Sequence	MASGSVAECLQQETTCPVCL QYFAEPMMLDCGHNICCAEL ARCWGTAETNVSCPQCRET PQRHMRPNRHLANVTQLVKQ LRTERPSGGGEMGVCEKHR EPLKLYCEEDQMPICVVCDR SREHRGHSVLPLEEAVEGFK EQIQQLDHLKRVKDLKRR RAQGEQARAELLSTQMERE KIVWEFEQLYHSLKEHEYRL
Specificity	Recombinant fusion protein containing a sequence corresponding to amino acids 1-200 of human TRIM27 (NP_006501.1).

ADDITIONAL INFORMATION


Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**



Immunohistochemistry analysis of TRIM27 in paraffin-embedded Mouse kidney using TRIM27 Rabbit polyclonal antibody (STJ118505) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of TRIM27 in paraffin-embedded Human liver using TRIM27 Rabbit polyclonal antibody (STJ118505) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of TRIM27 in paraffin-embedded Rat ovary using TRIM27 Rabbit polyclonal antibody (STJ118505) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.