# Anti-NAF1 antibody (150-400) (STJ119836)

## GENERAL INFORMATION

| | |
|---|---|
| Product Type | Primary antibodies |
| Applications | WB/ELISA |
| Host / Source | Rabbit |
| Reactivity | Human/Mouse/Rat |

## PRODUCT PROPERTIES


| | |
|---|---|
| Clonality | Polyclonal |
| Concentration | Lot specific |
| Conjugation | Unconjugated |
| Purification | Affinity purification |
| Dilution Range | WB:1:500-1:2000<br>ELISA:Recommended starting concentration is 1 Mu g/mL. Please optimize the concentration based on your specific assay requirements. |
| Formulation | PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3. |
| Isotype | IgG |
| Molecular Weight | Protein Mw: 54kDa<br>Observed Mw: 62-70kDa |
| Storage Instruction | Store at-20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles. |

## TARGET INFORMATION

| | |
|---|---|
| Gene ID | 92345 |
| Gene Symbol | NAF1 |
| UniProt ID | NAF1_HUMAN |
| Immunogen Region | 150-400 |
| Immunogen Sequence | PPVLSDGDDDLQIEKENKNF PLKTKDELLLNELPSVEELT IILPEDIELKPLGMVSSIIE QLVIIESMTNLPPVNEETVI FKSDRQAAGKIFEIFGPVAH PFYVLRFNSSDHIESKGIKI KETMYFAPSMKDFTQYIFTE KLKQDKGSDASWKNDQEPPP EALDFSDDEKEKEAKQRKKS QIQGRKKLKSEFNEPGEDFT EVHQNWNAHSSASEHAKGYR NREFTRGFSRARYPRSCHG |
| Specificity | Recombinant fusion protein containing a sequence corresponding to amino acids 150-400 of human NAF1 (NP_612395.2). |

## ADDITIONAL INFORMATION

| | |
|---|---|
| Note | STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS. |




Western blot analysis of various lysates using NAF1 Rabbit polyclonal antibody (STJ119836) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 Mu g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 60s.