

Anti-MT-ND1 antibody (1-100) (STJ119946)

GENERAL INFORMATION

Product Type	Primary antibodies
Applications	WB/IHC-P/IF/ICC/ELISA
Host / Source	Rabbit
Reactivity	Human/Mouse/Rat

PRODUCT PROPERTIES


Clonality	Polyclonal
Concentration	Lot specific
Conjugation	Unconjugated
Purification	Affinity purification
Dilution Range	WB:1:500-1:2000 IHC-P:1:50-1:200 IF/ICC:1:50-1:200 ELISA:Recommended starting concentration is 1 μ g/mL. Please optimize the concentration based on your specific assay requirements.
Formulation	PBS with 0.01% Thimerosal, 50% Glycerol, pH 7.3.
Isotype	IgG
Molecular Weight	Protein Mw: 36kDa Observed Mw: 36kDa
Storage Instruction	Store at -20°C for up to 1 year from the date of receipt, and avoid repeat freeze-thaw cycles.

TARGET INFORMATION


Gene ID	4535
Gene Symbol	MT-ND1
UniProt ID	NU1M_HUMAN
Immunogen Region	1-100
Immunogen Sequence	MPMANLLLLIVPIIAMAFLLTERKILGYMQLRKGPNNV GPYGLLQPFADAMKLFKEP LKPATSTITLYITAPTLALT IALLWTPPLPMPNPLVNLNL
Specificity	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human MT-ND1 (YP_003024026.1).

ADDITIONAL INFORMATION


Note **STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS.**




Immunofluorescence analysis of L929 cells using MT-ND1 Rabbit polyclonal antibody (STJ119946) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunofluorescence analysis of C6 cells using MT-ND1 Rabbit polyclonal antibody (STJ119946) at dilution of 1:100 (40x lens). Secondary antibody: Cy3 Goat Anti-Rabbit IgG (H+L) at 1:500 dilution. Blue: DAPI for nuclear staining.




Immunohistochemistry analysis of MT-ND1 in paraffin-embedded mouse kidney using MT-ND1 Rabbit polyclonal antibody (STJ119946) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of MT-ND1 in paraffin-embedded human esophageal using MT-ND1 Rabbit polyclonal antibody (STJ119946) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Immunohistochemistry analysis of MT-ND1 in paraffin-embedded rat ovary using MT-ND1 Rabbit polyclonal antibody (STJ119946) at dilution of 1:100 (40x lens). Perform microwave antigen retrieval with 10 mM PBS buffer pH 7.2 before commencing with immunohistochemistry staining protocol.



Western blot analysis of various lysates using MT-ND1 Rabbit polyclonal antibody (STJ119946) at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) (STJS000856) at 1:10000 dilution. Lysates/proteins: 25 μ g per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.