# Anti-Antistasin antibody (110-160 aa) (STJ13100712)

| GENERAL INFORMATION | |
|---|---|
| Product Type | Primary antibodies |
| Applications | IHC/WB |
| Host / Source | Rabbit |
| Reactivity | Hydra |

| PRODUCT PROPERTIES | |
|---|---|
| Clonality | Polyclonal |
| Conjugation | Unconjugated |
| Purification | IgG purified |
| Dilution Range | IHC, WB (confirmed by recombinant protein). A dilution of 1:300 to 1:2000 is recommended. The optimal dilution should be determined by the end user. Not yet tested in other applications. |
| Formulation | Lyophilised |
| Isotype | IgG |
| Storage Instruction | Maintain the lyophilised/reconstituted antibodies frozen at-20°C for long term storage and refrigerated at 2-8°C for a shorter term. When reconstituting, Glycerol (1:1) may be added for an additional stability. Avoid freeze and thaw cycles. |

| TARGET INFORMATION | |
|---|---|
| Immunogen Region | 110-160 aa |
| Specificity | Specific for Antistasin. |

| ADDITIONAL INFORMATION | |
|---|---|
| Note | STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS. |