

## Anti-c-Myc-Tag antibody (410-419 aa) (STJ13101366)

### GENERAL INFORMATION

|               |                    |
|---------------|--------------------|
| Product Type  | Primary antibodies |
| Applications  | IHC/WB             |
| Host / Source | Rabbit             |
| Reactivity    | cMyc-Tag           |

### PRODUCT PROPERTIES

|                     |   |
|---------------------|---|
| Clonality           | Polyclonal  |
| Conjugation         | Unconjugated  |
| Purification        | IgG purified  |
| Dilution Range      | IHC, WB using recombinant protein expressing {EQKLISEEDL (410-419) }. It is not recommended for tissue lysates. A dilution of 1:300 to 1:2000 is recommended. The optimal dilution should be determined by the end user.                          |
| Formulation         | Lyophilised   |
| Isotype             | IgG   |
| Storage Instruction | Maintain the lyophilised/reconstituted antibodies frozen at -20°C for long term storage and refrigerated at 2-8°C for a shorter term. When reconstituting, Glycerol (1:1) may be added for an additional stability. Avoid freeze and thaw cycles. |

### TARGET INFORMATION

|                  |                     |
|------------------|---------------------|
| Immunogen Region | 410-419 aa          |
| Specificity      | Specific for c-Myc. |

### ADDITIONAL INFORMATION

|      |   |
|------|---|
| Note | STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS. |
|------|---|