# Anti-NOGO antibody (460-510 aa) (STJ13101449)

## GENERAL INFORMATION

| | |
|---|---|
| Product Type | Primary antibodies |
| Applications | IHC/WB |
| Host / Source | Rabbit |
| Reactivity | Human/Rat/Mouse |

## PRODUCT PROPERTIES

| | |
|---|---|
| Clonality | Polyclonal |
| Conjugation | Unconjugated |
| Purification | IgG purified |
| Dilution Range | IHC, WB. A concentration of 10-50 μg/ml is recommended. The optimal concentration should be determined by the end user. Not yet tested in other applications. |
| Formulation | Lyophilised |
| Isotype | IgG |
| Storage Instruction | Maintain the lyophilised/reconstituted antibodies frozen at-20°C for long term storage and refrigerated at 2-8°C for a shorter term. When reconstituting, Glycerol (1:1) may be added for an additional stability. Avoid freeze and thaw cycles. |

## TARGET INFORMATION

| | |
|---|---|
| Immunogen Region | 460-510 aa |
| Specificity | Specific for Nogo. |

## ADDITIONAL INFORMATION

| | |
|---|---|
| Note | STRICTLY FOR FURTHER SCIENTIFIC RESEARCH USE ONLY (RUO). MUST NOT TO BE USED IN DIAGNOSTIC OR THERAPEUTIC APPLICATIONS. |